

Natural Pharmacy Business May 2017 references

CPD Fact File: It's all in the gut, by Stephen O'Hara

1. World Health Organization Factsheet 311 (2016):
<http://www.who.int/mediacentre/factsheets/fs311/en/>
2. World Obesity Federation: <http://www.worldobesity.org>
3. Public Health England:
https://www.noo.org.uk/NOO_about_obesity/obesity_and_health
4. Behan, D; Cox S; The Economic cost of Obesity. *Insurance Journal Jan.* 2011
5. Dombrowski SU, Avenell A, Sniehotta FF Long term maintenance of weight loss with non-surgical interventions in obese adults: systematic review and meta-analyses of randomised controlled trials *BMJ* 2014;348:g2646
6. Wing RR, Phelan S. Long-term weight loss maintenance. *Am J Clin Nutr* 2005;82:222-5
7. Avenell A, Broom J, Brown TJ, Poobalan A, Aucott L, Stearns SC, et al. Systematic review
8. of the long-term effects and economic consequences of treatments for obesity and implications for health improvement. *Health Technol Assess* 2004;8:iii-iv,1-182
9. Penn L, White M, Lindström J, den Boer AT, Blaak E, Eriksson JG, et al. Importance of
10. weight loss maintenance and risk prediction in the prevention of type 2 diabetes: analysis of European Diabetes Prevention Study RCT. *PLoS One* 2013;8:e57143.
11. Dombrowski SU, Avenell A, Sniehotta FF. Behavioural interventions for obese adults with
12. additional risk factors for morbidity: systematic review of effects on behaviour, weight and disease risk factors. *Obes Facts* 2010;3:377-96.
13. MacDougall, R., 2012. NIH Human Microbiome Project defines normal bacterial makeup of the body. Website: <http://www.nih.gov/news/health/jun2012/nhgri-13.htm>.
14. Nicholson, J.K., Holmes, E., Kinross, J., Burcelin, R., Gibson, G., Jia, W. and Pettersson, S., 2012. Host-gut microbiota metabolic interactions. *Science*, 336(6086), pp.1262-1267.
15. den Besten, G., van Eunen, K., Groen, A.K., Venema, K., Reijngoud, D.J. and Bakker, B.M., 2013. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Journal of Lipid Research*, 54(9), pp.2325-2340.
16. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, Semenkovich CF, Gordon JI: The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci USA* 2004;101:15718–15723
17. Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD, Gordon JI: Obesity alters gut microbial ecology. *Proc Natl Acad Sci USA* 2005;102:11070–11075
18. Perry, R.J., Peng, L., Barry, N.A., Cline, G.W., Zhang, D., Cardone, R.L., Petersen, K.F., Kibbey, R.G., Goodman, A.L. and Shulman, G.I., 2016. Acetate mediates a microbiome–brain– β -cell axis to promote metabolic syndrome. *Nature*, 534(7606), pp.213-217.
19. Ridaura, V.K., Faith, J.J., Rey, F.E., Cheng, J., Duncan, A.E., Kau, A.L., Griffin, N.W., Lombard, V., Henrissat, B., Bain, J.R. and Muehlbauer, M.J., 2013. Gut microbiota from twins discordant for obesity modulate metabolism in mice. *Science*, 341(6150), p.1241214.
20. Gough, E., Shaikh, H. and Manges, A.R., 2011. Systematic review of intestinal microbiota transplantation (fecal bacteriotherapy) for recurrent *Clostridium difficile* infection. *Clinical Infectious Diseases*, 53(10), pp.994-1002.

21. Alang, N. and Kelly, C.R., 2015, January. Weight gain after fecal microbiota transplantation. In *Open Forum Infectious Diseases* (Vol. 2, No. 1, p. ofv004). Oxford University Press.

Sugar control, by Mike Wakeman

1. Nahin RL, Byrd-Clark D, Stussman BJ, Kalyanaraman N. Disease severity is associated with the use of complementary medicine to treat or manage type-2 diabetes: data from the 2002 and 2007 National Health Interview Survey. *BMC Complement Altern Med*. 2012;12:193
2. Yeh GY, Eisenberg DM, Davis RB, Phillips RS. Use of complementary and alternative medicine among persons with diabetes mellitus: Results of a national survey. *Am J Public Health* 2002;92:1648-52
3. Broca C, Manteghetti M, Gross R, et al. 4-Hydroxyisoleucine: effects of synthetic and natural analogues on insulin secretion. *Eur J Pharmacol* 2000;390:339-45.
4. Madar Z, Abel R, Samish S, Arad J. Glucose-lowering effect of fenugreek in non-insulin dependent diabetics. *Eur J Clin Nutr* 1988;42:51-4
5. Kuriyan, R., Rajendran, R., Bantwal, G., and Kurpad, A. V. Effect of supplementation of *Coccinia cordifolia* extract on newly detected diabetic patients. *Diabetes Care* 2008;31(2):216-220.
6. Khan A, Safdar M, Ali Khan M, et al. Cinnamon improves glucose and lipids of people with type 2 diabetes. *Diabetes Care* 2003;26:3215-8.
7. Anderson RA, Broadhurst CL, Polansky MM, et al. Isolation and Characterization of Polyphenol Type-A Polymers from Cinnamon with Insulin-like Biological Activity. *J Agric Food Chem* 2004;52:65-70
8. Mang, B., Wolters, M., Schmitt, B., Kelb, K., Lichtinghagen, R., Stichtenoth, D. O., and Hahn, A. Effects of a cinnamon extract on plasma glucose, HbA_{1c}, and serum lipids in diabetes mellitus type 2. *Eur J Clin Invest* 2006;36(5):340-344
9. Vincent JB. The biochemistry of chromium. *J Nutr* 2000;130:715-8.
10. Lee NA, Reasner CA. Beneficial effect of chromium supplementation on serum triglyceride levels in NIDDM. *Diabetes Care* 1994;17:1449-52
11. Lima M, Cruz T, Pousada JC, et al. The effect of magnesium supplementation in increasing doses on the control of type 2 diabetes. *Diabetes Care* 1998;21:682-6.
12. Meyer KA, Kushi LH, Jacobs DR, et al. Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr* 2000;71:921-30
13. Rodriguez-Moran M, Guerrero-Romero F. Oral magnesium supplementation improves insulin sensitivity and metabolic control in type 2 diabetic subjects: A randomized double-blind controlled trial. *Diabetes Care* 2003;26:1147-52
14. Rude RK. Magnesium deficiency: a cause of heterogeneous disease in humans. *J Bone Miner Res* 1998;13:749-58
15. Vuksan V, Stavro MP, Sievenpiper JL, et al. Similar postprandial glycemic reductions with escalation of dose and administration time of American ginseng in type 2 diabetes. *Diabetes Care* 2000;23:1221-6.
16. Sievenpiper JL, Arnason JT, Leiter LA, Vuksan V. Decreasing, null and increasing effects of eight popular types of ginseng on acute postprandial glycemic indices in healthy humans: the role of ginsenosides. *J Am Coll Nutr* 2004;23:248-58.

17. Xia, X., Yan, J., Shen, Y., Tang, K., Yin, J., Zhang, Y., Yang, D., Liang, H., Ye, J., and Weng, J. Berberine improves glucose metabolism in diabetic rats by inhibition of hepatic gluconeogenesis. *PLoS.One*. 2011;6(2):e16556.
18. Lu, S. S., Yu, Y. L., Zhu, H. J., Liu, X. D., Liu, L., Liu, Y. W., Wang, P., Xie, L., and Wang, G. J. Berberine promotes glucagon-like peptide-1 (7-36) amide secretion in streptozotocin-induced diabetic rats. *J Endocrinol*. 2009;200(2):159-165.
19. Zhang Y, Li X, Zou D, et al. Treatment of type 2 diabetes and dyslipidemia with the natural plant alkaloid berberine. *J Clin Endocrinol Metab* 2008;93:2559-65.
20. Yin, J., Xing, H., and Ye, J. Efficacy of berberine in patients with type 2 diabetes mellitus. *Metabolism* 2008;57(5):712-717
21. Boivin M, Flourie B, Rizza RA, Go VL, DiMagno EP. Gastrointestinal and metabolic effects of amylase inhibition in diabetics. *Gastroenterology*. 1988;94(2):387-94
22. Anderson JW, Allgood LD, Turner J, et al. Effects of psyllium on glucose and serum lipid responses in men with type 2 diabetes and hypercholesterolemia. *Am J Clin Nutr* 1999;70:466-73
23. Russo A, Stevens JE, Wilson T, et al. Guar attenuates fall in postprandial blood pressure and slows gastric emptying of oral glucose in type 2 diabetes. *Dig Dis Sci* 2003;48:1221-9.
24. Jenkins DJ, Wesson V, Wolever TM, et al. Wholemeal versus wholegrain breads: proportion of whole or cracked grain and the glycaemic response. *BMJ* 1988;297:958-60
25. Pick ME, Hawrysh ZJ, Gee MI, et al. Oat bran concentrate bread products improve long-term control of diabetes: a pilot study. *J Am Diet Assoc* 1996;96:1254-61
26. Akilen, R., Tsiami, A., Devendra, D., and Robinson, N. Glycated haemoglobin and blood pressure-lowering effect of cinnamon in multi-ethnic Type 2 diabetic patients in the UK: a randomized, placebo-controlled, double-blind clinical trial. *Diabet Med*. 2010;27(10):1159-1167.
27. Mang, B., Wolters, M., Schmitt, B., Kelb, K., Lichtinghagen, R., Stichtenoth, D. O., and Hahn, A. Effects of a cinnamon extract on plasma glucose, HbA, and serum lipids in diabetes mellitus type 2. *Eur J Clin Invest* 2006;36(5):340-344
28. Doi K, Matsuura M, Kawara A, Baba S. Treatment of diabetes with glucomannan (konjac mannan). *Lancet* 1979;1:987-8.
29. Cairella M, Marchini GAD. [Evaluation of the action of glucomannan on metabolic parameters and on the sensation of satiation in overweight and obese patients]. [Article in Italian] *Clin Ter* 1995;146:269-74
30. Konrad T, Vicini P, Kusterer K, et al. Alpha-lipoic acid treatment decreases serum lactate and pyruvate concentrations and improves glucose effectiveness in lean and obese patients with Type 2 diabetes. *Diabetes Care* 1999;22:280-7
31. Ziegler D, Hanefeld M, Ruhnau K, et al. Treatment of symptomatic diabetic polyneuropathy with the antioxidant alpha-lipoic acid: A 7-month, multicenter, randomized, controlled trial (ALADIN III Study). *Diabetes Care* 1999;22:1296-301
32. Hodgson JM, Watts GF, Playford DA, et al. Coenzyme Q10 improves blood pressure and glycaemic control: a controlled trial in subjects with type 2 diabetes. *Eur J Clin Nutr* 2002;56:1137-42
33. Henriksen JE, Andersen CB, Hother-Nielsen O, et al. Impact of ubiquinone (coenzyme Q10) treatment on glycaemic control, insulin requirement and well-being in patients with Type 1 diabetes mellitus. *Diabet Med* 1999;16:312-8
34. Hernandez-Ojeda, J., Cardona-Munoz, E. G., Roman-Pintos, L. M., Troyo-Sanroman, R., Ortiz-Lazareno, P. C., Cardenas-Meza, M. A., Pascoe-Gonzalez, S., and Miranda-Diaz, A.

- G. The effect of ubiquinone in diabetic polyneuropathy: a randomized double-blind placebo-controlled study. *J Diabetes Complications* 2012;26(4):352-358.
35. Vuksan V, Whitham D, Sievenpiper JL, et al. Supplementation of conventional therapy with the novel grain Salba (*Salvia hispanica* L.) improves major and emerging cardiovascular risk factors in type 2 diabetes: results of a randomized controlled trial. *Diabetes Care* 2007;30:2804-10.